

**THIS PAGE IS INSERTED BY OIPE SCANNING
AND IS NOT PART OF THE OFFICIAL RECORD**

Best Available Images

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

BLACK BORDERS

TEXT CUT OFF AT TOP, BOTTOM OR SIDES

FADED TEXT

BLURRY OR ILLEGIBLE TEXT

SKEWED/SLANTED IMAGES

COLORED PHOTOS HAVE BEEN RENDERED INTO BLACK AND WHITE

VERY DARK BLACK AND WHITE PHOTOS

UNDECIPHERABLE GRAY SCALE DOCUMENTS

**IMAGES ARE THE BEST AVAILABLE
COPY. AS RESCANNING *WILL NOT*
CORRECT IMAGES, PLEASE DO NOT
REPORT THE IMAGES TO THE
PROBLEM IMAGE BOX.**

Figure 1 Human T1R2/T1R3 functions as a sweet taste receptor




Figure 2 T1R2 may control T1R2/T1R3 ligand specificity





Figure 3 Human T1R1/T1R3 functions as an umami taste receptor


a




b



c



A.



H2-TT-Myc

H2-PDZIP-TT-Myc

B.




Figure 4 PDZIP facilitate the surface expression of human T1R2.

- A. Immunofluorescence staining of Myc-tagged hT1R2 indicates that PDZIP significantly increases the amount of human T1R2 protein on the plasma membrane (Staszewski, May 15th, 2001, Notebook No. 2, page 76-77).
- B. FACS analysis data demonstrating the same result (Staszewski, June 4th, 2001, Notebook No. 2, page 79). Myc-tagged human T1R2: Green line. Myc-tagged human T1R2 with PDZIP: black line.

Figure 5




Figure 6

